

## County Minimum Urinalysis Testing Requirements

1. Pregnant and Parenting Women Programs:
  - a. Minimum testing needs for Pregnant and Parenting Women Programs are for the following four-panel screen plus adulterants including creatinine:
    - (1) Amphetamine/methamphetamine - 1000 ng/mL
    - (2) Cocaine - 300 ng/mL
    - (3) Opiates - 300 ng/LI
    - (4) Cannabinoid (THC) at 50 ng/mL; all positive tests quantitated
  - b. All positive tests must be confirmed by Gas Chromatography - Mass Spectroscopy (GC/MS) or Liquid Chromatography/Tandem Mass Spectroscopy (LCMSMS).
  - c. Testing Confirmation Cutoffs:
    - (1) Amphetamine/Methamphetamine - 500 ng/mL
    - (2) MDMA/MDA - 250 ng/mL
    - (3) Cocaine met. (Benzoylecgonine) - 150 ng/mL
    - (4) Opiates testing include the following drugs – 300 ng/ml:
      - (a) Morphine
      - (b) Codeine
      - (c) Hydrocodone
      - (d) Hydromorphone
      - (e) Oxycodone
  - d. THC testing include the following drug:
    - (1) Carboxy-THC – 15 ng/mL
2. Opiate Substitution Treatment Programs (OSTP)
  - a. Minimum testing needs for OSTP require the following five-panel screens plus adulterants including Creatinine.

- (1) Amphetamine/methamphetamine - 1000 ng/mL
  - (2) Benzodiazepines - 200 ng/mL
  - (3) Cocaine - 300 ng/mL
  - (4) Methadone 300 ng/mL
  - (5) Opiates - 300 ng/mL
- b. Cocaine, Methadone and Opiates must be included in the five-panel screen. Programs may substitute up to two other drugs, including THC, in the five-panel screen for documented clinical purposes. Confirmation Cutoffs:
- (1) Amphetamine/Methamphetamine – 500 ng/ml
  - (2) MDMA – 250 ng/mL
  - (3) Benzodiazepines – 100 ng/ml
    - (a) Nordiazepam
    - (b) Oxazepam
    - (c) Temezepam
    - (d) Lorazepam
    - (e) Clonazepam
    - (f) Aprazolam
  - (4) Cocaine met. (Benzoylecgonine) – 150 ng/ml
  - (5) Methadone – 300 ng/ml
  - (6) EDDP – 100 ng/ml
  - (7) Opiates – 300 ng/ml
  - (8) Morphine
  - (9) Codeine
  - (10) Hydromorphone
  - (11) Oxycodone
- c. Opiate positive screens must be confirmed for codeine, morphine, hydrocodone, hydromorphone, oxycodone, and 6-acetylmorphine.

- d. Benzodiazepine positive screens must be confirmed for nordiazepam, oxazepam, temazepam, lorazepam, clonazepam, and alprazolam.
3. Outpatient and Residential Treatment programs are required to perform a minimum of a four-panel screen plus adulterants including creatinine, with established confirmation cutoff.
- a. Amphetamine/methamphetamine - 1000 ng/ml
  - b. Cocaine - 300 ng/ml
  - c. Opiates - 300 ng/ml
  - d. Cannabinoid (THC) at 50 ng/ml; all positive screens quantitated
  - e. All positive screens must be confirmed by GC/MS. Confirmation Cutoffs:
  - f. Amphetamine/Methamphetamine - 500 ng/ml
  - g. MDMA/MDA - 250 ng/ml
  - h. Cocaine met. (Benzoylecgonine) - 150 ng/ml
  - i. Opiates – 300 ng/ml:
    - (1) Morphine
    - (2) Codeine
    - (3) Hydrcodone
    - (4) Hydromorphone
    - (5) Oxycodone
  - j. THC:
    - (1) Carboxy-THC - 15 ng/ml
4. Alcohol testing should be part of the drug testing panel only when the donor is suspect by odor or overt behavior.